



BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood // Levine EMB Agar

111-252258-N-00 , April 2015

QUALITY CONTROL PROCEDURES

I. INTRODUCTION

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood (TSA5%SB) is used for the growth of fastidious organisms and for the visualization of hemolytic reactions. **Levine EMB Agar** is a slightly selective and differential medium for the isolation, cultivation and differentiation of gram-negative microorganisms isolated from both clinical and nonclinical specimens. It is widely used for the examination of materials of sanitary importance for the presence of coliforms.

II. PERFORMANCE TEST PROCEDURE

A. **BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood**

- Inoculate representative samples with dilutions of the cultures listed below.
 - Using a volumetric pipettor or equivalent method, deliver 0.1 mL of a dilution yielding 30–300 CFU to each plate and spread inoculate using a sterile glass spreader.
 - Incubate the *Staphylococcus aureus* and *Escherichia coli* strains at $35 \pm 2^\circ\text{C}$ in an aerobic atmosphere and the *Streptococcus* strains at $35 \pm 2^\circ\text{C}$ in an aerobic atmosphere supplemented with 3–5% carbon dioxide.
- Examine plates after 18–24 h for growth, colony size and hemolytic reactions.
- Expected Results

CLSI Organisms	ATCC®	Recovery
* <i>Streptococcus pyogenes</i>	19615	Growth, beta hemolysis
* <i>Streptococcus pneumoniae</i>	6305	Growth, alpha hemolysis
* <i>Staphylococcus aureus</i>	25923	Growth
* <i>Escherichia coli</i>	25922	Growth

*Recommended organism strain for User Quality Control.

B. **Levine EMB Agar**

- Inoculate representative samples with dilutions of the cultures listed below.
 - Streak the plates for isolation using cultures diluted to yield 10^3 – 10^5 CFU/plate.
 - Incubate plates at $35 \pm 2^\circ\text{C}$ in an aerobic atmosphere.
 - Include **BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood** plates as nonselective controls for all organisms.
- Examine plates after 18–24 h for amount of growth, colony size, pigmentation and selectivity.
- Expected Results

CLSI Organisms	ATCC®	Recovery
* <i>Escherichia coli</i>	25922	Growth, blue-black colonies, with partial green metallic sheen
* <i>Salmonella enterica</i> subsp. <i>enterica</i> serovar Typhimurium	14028	Growth, colorless to amber colonies
* <i>Enterococcus faecalis</i>	29212	Inhibition (partial)
Additional Organism		
<i>Shigella flexneri</i>	12022	Growth, colorless to amber

*Recommended organism strain for User Quality Control.

III. ADDITIONAL QUALITY CONTROL

1. Examine plates as described under "Product Deterioration."
2. Visually examine representative plates to assure that any existing physical defects will not interfere with use.
3. Determine the pH potentiometrically at room temperature for adherence to the specification of 7.4 ± 0.2 (TSA5%SB) and 7.3 ± 0.2 (Levine EMB Agar).
4. Note the firmness of plates during the inoculation procedure.
5. Incubate uninoculated representative plates aerobically at $35 \pm 2^\circ\text{C}$ for 72 h and examine for microbial contamination.

PRODUCT INFORMATION

IV. INTENDED USE

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood is used for cultivating fastidious microorganisms and for the visualization of hemolytic reactions produced by many bacterial species.

Levine EMB Agar is a selective and differential plating medium for the isolation of gram-negative enteric bacteria.

V. SUMMARY AND EXPLANATION

A. **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood

The nutritional composition of **BD BBL™ Trypticase™** Soy Agar has made it a popular medium, both unsupplemented and as a base for media containing blood. **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood is extensively used for the recovery and cultivation of fastidious microbial species and for the determination of hemolytic reactions which are important differentiating characteristics for bacteria, especially *Streptococcus* species.

B. Levine EMB Agar

Shortly following the publication by Holt-Harris and Teague of a paper describing a new culture medium for the differentiation of enteric microorganisms through the use of eosin and methylene blue dyes,¹ Levine described a modification of their formulation which he claimed gave better differentiation between what are now referred to as *Escherichia* and *Enterobacter* species.² The two formulations differ in that Levine EMB Agar does not contain sucrose. Both of these formulations were developed to improve upon the differentiating properties of Endo Agar,³ which was developed previously. Levine EMB Agar utilizes dyes as selective agents. It is listed for use in the microbiological examination of dairy products and foods by the American Public Health Association.^{4,5}

VI. PRINCIPLES OF THE PROCEDURE

A. **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood

The combination of casein and soy peptones in the **BD BBL™ Trypticase™** Soy Agar base render the medium highly nutritious by supplying organic nitrogen, particularly amino acids and larger-chained peptides. The sodium chloride maintains osmotic equilibrium. Defibrinated sheep blood is the most widely used blood for enriching agar base media.⁶ Hemolytic reactions of streptococci are proper and growth of *Haemophilus hemolyticus*, a nonpathogen whose hemolytic colonies are indistinguishable from those of betahemolytic streptococci, is inhibited.

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood provides excellent growth and beta hemolysis by *Streptococcus pyogenes* (Lancefield group A) and also provides excellent growth and appropriate hemolytic reactions with other fastidious organisms. It is suitable for use with low concentration (0.04 unit) bacitracin discs (**BD Taxo™ A**) for presumptive identification of group A streptococci (*S. pyogenes*).

B. Levine EMB Agar

The eosin Y and methylene blue dyes in Levine EMB Agar render the medium slightly selective in that they inhibit gram-positive bacteria to a limited degree. These dyes also play a role in differentiating between lactose fermenters and lactose-nonfermenters due to the presence or absence of dye uptake in the bacterial colonies. Coliforms, as lactose fermenting organisms, are visualized as blue-black colonies

whereas colonies of *Salmonella* and *Shigella*, as lactose-nonfermenters, appear colorless, transparent or amber in color. Some gram-positive bacteria, such as fecal streptococci, staphylococci and yeasts, will grow on this medium and usually form pinpoint colonies. A number of nonpathogenic, lactose-nonfermenting gram-negative bacteria will grow on this medium and must be distinguished from the pathogenic strains by additional biochemical tests.

VII. REAGENTS

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood

Approximate Formula* Per Liter Purified Water

Pancreatic Digest of Casein	14.5 g
Papaic Digest of Soybean Meal	5.0 g
Sodium Chloride	5.0 g
Agar.....	14.0 g
Growth Factors	1.5 g
Defibrinated Sheep Blood	5 %

*Adjusted and/or supplemented as required to meet performance criteria.

Levine EMB Agar

Approximate Formula* Per Liter Purified Water

Pancreatic Digest of Gelatin	10.0 g
Lactose	10.0 g
Dipotassium Phosphate	2.0 g
Eosin Y	0.4 g
Methylene Blue	0.065 g
Agar	15.0 g

*Adjusted and/or supplemented as required to meet performance criteria.

Warnings and Precautions:

For *in vitro* Diagnostic Use in Taiwan.

If excessive moisture is observed, invert the bottom over an off-set lid and allow to air dry in order to prevent formation of a seal between the top and bottom of the plate during incubation. Pathogenic microorganisms, including hepatitis viruses and Human Immunodeficiency Virus, may be present in clinical specimens. "Standard Precautions"⁷⁻¹⁰ and institutional guidelines should be followed in handling all items contaminated with blood and other body fluids. After use, prepared plates, specimen containers and other contaminated materials must be sterilized by autoclaving before discarding.

Storage Instructions: On receipt, store plates in the dark at 2–8°C. Avoid freezing and overheating. Do not open until ready to use. Minimize exposure to light. Prepared plates stored in their original sleeve wrapping at 2–8°C until just prior to use may be inoculated up to the expiration date and incubated for recommended incubation times. Allow the medium to warm to room temperature before inoculation.

Product Deterioration: Do not use plates if they show evidence of microbial contamination, discoloration, drying, cracking or other signs of deterioration.

VIII. specimen collection and handling

Specimens suitable for culture may be handled using various techniques. For detailed information, consult appropriate texts.^{11,12} Specimens should be obtained before antimicrobial therapy has been administered. Provision must be made for prompt delivery to the laboratory.

IX. PROCEDURE

Material Provided: BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood //Levine EMB Agar

Materials Required But Not Provided: Ancillary culture media, reagents, quality control organisms and laboratory equipment as required.

Test Procedure: Observe aseptic techniques.

The agar surface should be smooth and moist, but without excessive moisture. Streak the specimen as soon as possible after it is received in the laboratory. The streak plate is used primarily to isolate pure cultures from specimens containing mixed flora. Alternatively, if material is being cultured directly from a swab, roll the swab over a small area of the surface at the edge; then streak from this inoculated area. Incubate plates, protected from light, at $35 \pm 2^\circ\text{C}$ for 18–24 h in an aerobic or CO_2 -enriched atmosphere, depending on the source of the specimen or the microorganisms suspected of being present.

User Quality Control: See "Quality Control Procedures."

Quality Control requirements must be performed in accordance with applicable local, state and/or federal regulations or accreditation requirements and your laboratory's standard Quality Control procedures. It is recommended that the user refer to pertinent CLSI (formerly NCCLS) guidance and CLIA regulations for appropriate Quality Control practices.

X. RESULTS

After incubation most plates will show an area of confluent growth. Because the streaking procedure is, in effect, a "dilution" technique, diminishing numbers of microorganisms are deposited on the streaked areas. Consequently, one or more of these areas should exhibit isolated colonies of the organisms contained in the specimen.

Typical results on **BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood** are as follows:

1. Hemolytic streptococci may appear as translucent or opaque, grayish, small (1 mm), or large matte and mucoid (2–4 mm) colonies, encircled by a zone of hemolysis. Gram stains should be made and examined to check the macroscopic findings. (Other organisms which may cause hemolysis include *Listeria*, various corynebacteria, hemolytic staphylococci, *Escherichia coli* and *Pseudomonas*.) In reporting, approximate quantitation of the number of colonies of hemolytic streptococci may be helpful to the clinician.
2. Pneumococci usually appear as very flat, smooth, translucent, grayish and sometimes mucoid colonies surrounded by a narrow zone of "green" (alpha) hemolysis.
3. Staphylococci appear as opaque, white to gold-yellow colonies with or without zones of beta hemolysis.
4. *Listeria*. Small zones of beta hemolysis are produced. They may be distinguished by their rod shape in stains, and by motility at room temperature.
5. Other organisms representing minimal flora and clinically significant isolates can also be expected to grow on this nonselective formulation.

Typical colonial morphology on Levine EMB Agar is as follows:

<i>E. coli</i>	Large, blue-black, with partial green metallic sheen
<i>Enterobacter/Klebsiella</i>	Large, mucoid, blue-black
<i>Proteus</i>	Large, colorless
<i>Salmonella</i>	Large, colorless
<i>Shigella</i>	Medium to Large, colorless
<i>Pseudomonas</i>	Irregular, colorless
Gram-positive bacteria.....	No growth to slight growth

XI. limitations of the procedure

For identification, organisms must be in pure culture. Morphological, biochemical, and/or serological tests should be performed for final identification. Consult appropriate texts for detailed information and recommended procedures.¹¹⁻¹⁶

XII. performance Characteristics

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood

BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood was used as a control in a study using broth-enhanced culture (Todd Hewitt) and Optical Immunoassay method for the diagnosis of hemolytic streptococcal infection. Five hundred two (502) specimens were tested. TSA with 5% Sheep Blood had a

sensitivity and specificity of 92.5% and 99.4%, respectively.¹⁷ Nguyen et al. used **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood as the “gold standard” for the detection of group B *Streptococcus* from the lower genital tract of pregnant women.¹⁸ In another study, Rossmann et al. successfully reisolated *Lautropia mirabilis* on **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood from the oral cavities of human immunodeficiency virus infected children.¹⁹ Of the 85 children evaluated in this study, 35 (41.4%) were positive for *L. mirabilis*. Isenberg et al. used **BD BBL™ Trypticase™** Soy Agar with 5% Sheep Blood as a control to evaluate the recovery of *Enterococcus* from a selective medium under study.²⁰ Two hundred fifty (250) group D streptococcal strains isolated from clinical material and 8 strains obtained from the National Communicable Disease Center (Atlanta, Ga.) were used.

XIII. availability

Cat. No.	Description
252258	BD BBL™ Trypticase™ Soy Agar with 5% Sheep Blood //Levine EMB Agar, Ctn. of 100 plates

XIV. REFERENCES

- Holt-Harris, J.E., and O. Teague. 1916. A new culture medium for the isolation of *Bacillus typhosus* from stools. J. Inf. Dis. 18:596-600.
- Levine, M. 1918. Differentiation of *B. coli* and *B. aerogenes* on a simplified eosin-methylene blue agar. J. Inf. Dis. 23:43-47.
- Endo, S. 1904. Ueber ein verfahren zum nachweis der typhusbacillen. Centr. f. Bakt. 35:109-110.
- Wehr, H.M. and J.H. Frank (ed.). 2004. Standard methods for the examination of dairy products, 17th ed. American Public Health Association. Washington, D.C.
- Downes, F.P. and K. Ito (ed.). 2001. Compendium of methods for the microbiological examination of foods, 4th ed. American Public Health Association, Washington, D.C.
- Vera, H.D., and D.A. Power. 1980. Culture media, p. 969. In E.H. Lennette, A. Balows, W.J. Hausler, Jr., and J.P. Truant (ed.), Manual of clinical microbiology, 3rd ed. American Society for Microbiology, Washington, D.C.
- Clinical and Laboratory Standards Institute. 2014. oved Guideline M29-A4 Protection of laboratory workers from occupationally acquired infections, 4th ed. CLSI, Wayne, PA.
- Garner, J.S. 1996. Hospital Infection Control Practices Advisory Committee, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Guideline for isolation precautions in hospitals. Infect. Control Hospital Epidemiol. 17:53-80.
- U.S. Department of Health and Human Services. 2007. Biosafety in microbiological and biomedical laboratories, HHS Publication (CDC), 5th ed. U.S. Government Printing Office, Washington, D.C.
- Directive 2000/54/EC of the European Parliament and of the Council of 18 September 2000 on the protection of workers from risks related to exposure to biological agents at work (seventh individual directive within the meaning of Article 16(1) of Directive 89/391/EEC). Official Journal L262,17/10/2000, p. 0021-0045.
- Murray, P.R., E.J. Baron, J.H. Jorgensen, M.A. Pfaller, and R.H. Tenover (ed.). 2011. Manual of clinical microbiology, 10th ed. American Society for Microbiology, Washington, D.C.
- Forbes, B.A., D.F. Sahm, and A.S. Weissfeld. 2013. Bailey and Scott's diagnostic microbiology, 13th ed. Mosby, Inc., St. Louis.
- Holt, J.G., N.R. Krieg, P.H.A. Sneath, J.T. Staley, and S.T. Williams (ed.). 1994. Bergey's Manual™ of determinative bacteriology, 9th ed. Williams & Wilkins, Baltimore.
- MacFaddin, J.F. 2000. Biochemical tests for identification of medical bacteria, 3rd ed. Lippincott Williams & Wilkins, Baltimore.
- Koneman, E.W., S.D. Allen, W.M. Janda, P.C. Schreckenberger, and W.C. Winn, Jr. 2004. Color atlas and textbook of diagnostic microbiology, 6th ed. Lippincott-Raven, Philadelphia.
- Isenberg, H.D. (ed.). 2010. Clinical microbiology procedures handbook, vol. 1, 2 and 3, 3rd ed. American Society for Microbiology, Washington, D.C.
- Fries, S.M. 1995. Diagnosis of group A streptococcal pharyngitis in a private clinic: comparative evaluation of an optical immunoassay method and culture. J. Pediatr. 126:933-936.
- Nguyen, T.M., et al. 1998. Detection of group B streptococcus: comparison of an optical immunoassay with direct plating and broth-enhanced culture methods. J. Matern. Fetal. Med. 7:172-176.
- Rossmann, S.N. et al. 1998. Isolation of *Lautropia mirabilis* from oral cavities of human immunodeficiency virus-infected children. J. Clin. Microbiol. 36:1756-1760.
- Isenberg, H.D., D. Goldberg, and J. Sampson, 1970. Laboratory studies with a selective medium. Appl. Microbiol. 20:433-436.

XV. FURTHER INFORMATION

For further information please contact your local BD representative.

Nippon Becton Dickinson Company, Ltd.

1 Aza Gotanda, Tsuchifune, Fukushima City, Fukushima, Japan

e-mail : BD-eDial@bd.com

WEB : <http://www.bd.com/jp/>

ATCC is a trademark of the American Type Culture Collection.

BD, BD Logo, and all other trademarks are property of Becton, Dickinson and Company. ©2015 BD.